

# The spectrum of simplicial volume of non-compact manifolds

Nicolaus Heuer, Clara Löh

October 24, 2020

## Abstract

We show that, in dimension at least 4, the set of locally finite simplicial volumes of oriented connected open manifolds is  $[0, \infty]$ . Moreover, we consider the case of tame open manifolds and some low-dimensional examples.

## 1 Introduction

Simplicial volumes are invariants of manifolds defined in terms of the  $\ell^1$ -semi-norm on singular homology [Gro82].

**Definition 1.1** (simplicial volume). Let  $M$  be an oriented connected  $d$ -manifold without boundary. Then the *simplicial volume of  $M$*  is defined by

$$\|M\|^{\text{lf}} := \inf\{|c|_1 \mid c \in C_d^{\text{lf}}(M; \mathbb{R}) \text{ is a fundamental cycle of } M\},$$

where  $C_*^{\text{lf}}$  denotes the locally finite singular chain complex. If  $M$  is compact, then we also write  $\|M\| := \|M\|^{\text{lf}}$ . Using relative fundamental cycles, the notion of simplicial volume can be extended to oriented manifolds with boundary.

Simplicial volumes are related to negative curvature, volume estimates, and amenability [Gro82]. In the present article, we focus on simplicial volumes of *non-compact* manifolds. Only few concrete results are known in this context: There are computations for certain locally symmetric spaces [LS09a, LS09b, BKK14, KK15] as well as the general volume estimates [Gro82], vanishing results [Gro82, FM18], and finiteness results [Gro82, Löh08].

Let  $d \in \mathbb{N}$ , let  $M(d)$  be the class of all oriented closed connected  $d$ -manifolds, and let  $M^{\text{lf}}(d)$  be the class of all oriented connected manifolds without boundary. Then we set  $\text{SV}(d) := \{\|M\| \mid M \in M(d)\}$  and

$$\text{SV}^{\text{lf}}(d) := \{\|M\|^{\text{lf}} \mid M \in M^{\text{lf}}(d)\}.$$

It is known that  $\text{SV}(d)$  is countable and that this set has no gap at 0 if  $d \geq 4$ :

**Theorem 1.2** ([HL20, Theorem A]). *Let  $d \in \mathbb{N}_{\geq 4}$ . Then  $\text{SV}(d)$  is dense in  $\mathbb{R}_{\geq 0}$  and  $0 \in \text{SV}(d)$ .*

---

© N. Heuer, C. Löh 2020. This work was supported by the CRC 1085 *Higher Invariants* (Universität Regensburg, funded by the DFG).  
MSC 2010 classification: 57N65

In contrast, if we allow non-compact manifolds, we can realise *all* non-negative real numbers:

**Theorem A.** *Let  $d \in \mathbb{N}_{\geq 4}$ . Then  $\text{SV}^{\text{lf}}(d) = [0, \infty]$ .*

The proof uses the no-gap theorem Theorem 1.2 and a suitable connected sum construction.

If we restrict to tame manifolds, then we are in a similar situation as in the closed case:

**Theorem B.** *Let  $d \in \mathbb{N}$ . Then the set  $\text{SV}_{\text{tame}}^{\text{lf}}(d) \subset [0, \infty]$  is countable. In particular, the set  $[0, \infty] \setminus \text{SV}_{\text{tame}}^{\text{lf}}(d)$  is uncountable.*

As an explicit example, we compute  $\text{SV}^{\text{lf}}(2)$  and  $\text{SV}_{\text{tame}}^{\text{lf}}(2)$  (Proposition 4.2) as well as  $\text{SV}_{\text{tame}}^{\text{lf}}(3)$  (Proposition 4.3). The case of non-tame 3-manifolds seems to be fairly tricky.

**Question 1.3.** What is  $\text{SV}^{\text{lf}}(3)$  ?

As  $\text{SV}(4) \subset \text{SV}_{\text{tame}}^{\text{lf}}(4)$ , we know that  $\text{SV}_{\text{tame}}^{\text{lf}}(4)$  contains arbitrarily small transcendental numbers [HL19].

From a geometric point of view, the so-called Lipschitz simplicial volume is more suitable for Riemannian non-compact manifolds than the locally finite simplicial volume. It is therefore natural to ask the following:

**Question 1.4.** Do Theorem A and Theorem B also hold for the Lipschitz simplicial volume of oriented connected open Riemannian manifolds?

## Organisation of this article

Section 2 contains the proof of Theorem A. The proof of Theorem B is given in Section 3. The low-dimensional case is treated in Section 4.

## 2 Proof of Theorem A

Let  $d \in \mathbb{N}_{\geq 4}$  and let  $\alpha \in [0, \infty]$ . Because  $\text{SV}(d)$  is dense in  $\mathbb{R}_{\geq 0}$  (Theorem 1.2), there exists a sequence  $(\alpha_n)_{n \in \mathbb{N}}$  in  $\text{SV}(d)$  with  $\sum_{n=0}^{\infty} \alpha_n = \alpha$ .


### 2.1 Construction

We first describe the construction of a corresponding oriented connected open manifold  $M$ : For each  $n \in \mathbb{N}$ , we choose an oriented closed connected  $d$ -manifold  $M_n$  with  $\|M_n\| = \alpha_n$ . Moreover, for  $n > 0$ , we set

$$W_n := M_n \setminus (B_{n,-}^{\circ} \sqcup B_{n,+}^{\circ}),$$

where  $B_{n,-} = i_{n,-}(D^d)$  and  $B_{n,+} = i_{n,+}(D^d)$  are two disjointly embedded closed  $d$ -balls in  $M_n$ . Similarly, we set  $W_0 := M_0 \setminus B_{0,+}^{\circ}$ . Furthermore, we choose an orientation-reversing homeomorphism  $f_n: S^{d-1} \rightarrow S^{d-1}$ . We then consider the infinite ‘‘linear’’ connected sum manifold (Figure 1)

$$\begin{aligned} M &:= M_0 \# M_1 \# M_2 \# \dots \\ &= (W_0 \sqcup W_1 \sqcup W_n \sqcup \dots) / \sim, \end{aligned}$$

Figure 1: The construction of  $M$  for the proof of Theorem A

where  $\sim$  is the equivalence relation generated by

$$i_{n+1,-}(x) \sim i_{n,+}(f_n(x))$$

for all  $n \in \mathbb{N}$  and all  $x \in S^{d-1} \subset D^d$ ; we denote the induced inclusion  $W_n \rightarrow M$  by  $i_n$ . By construction,  $M$  is connected and inherits an orientation from the  $M_n$ .

## 2.2 Computation of the simplicial volume

We will now verify that  $\|M\|^{\text{lf}} = \alpha$ :

**Claim 2.1.** *We have  $\|M\|^{\text{lf}} \leq \alpha$ .*

*Proof.* The proof is a straightforward adaption of the chain-level proof of sub-additivity of simplicial volume with respect to amenable glueings.

In particular, we will use the uniform boundary condition [MM85] and the equivalence theorem [Gro82, BBF<sup>+</sup>14]:

**UBC** The chain complex  $C_*(S^{d-1}; \mathbb{R})$  satisfies  $(d-1)$ -UBC, i.e., there is a constant  $K$  such that: For each  $c \in \text{im } \partial_d \subset C_{d-1}(S^{d-1}; \mathbb{R})$ , there exists a chain  $b \in C_d(S^{d-1}; \mathbb{R})$  with

$$\partial_d b = c \quad \text{and} \quad |b|_1 \leq K \cdot |c|_1.$$

**EQT** Let  $N$  be an oriented closed connected  $d$ -manifold, let  $B_1, \dots, B_k$  be disjointly embedded  $d$ -balls in  $N$ , and let  $W := N \setminus (B_1^\circ \cup \dots \cup B_k^\circ)$ . Moreover, let  $\epsilon \in \mathbb{R}_{>0}$ . Then

$$\|N\| = \inf \{ |z|_1 \mid z \in Z(W; \mathbb{R}), |\partial_d z|_1 \leq \epsilon \},$$

where  $Z(W; \mathbb{R}) \subset C_d(W; \mathbb{R})$  denotes the set of all relative fundamental cycles of  $W$ .

Let  $\epsilon \in \mathbb{R}_{>0}$ . By EQT, for each  $n \in \mathbb{N}$ , there exists a relative fundamental cycle  $z_n \in Z(W_n; \mathbb{R})$  with

$$|z_n|_1 \leq \alpha_n + \frac{1}{2^n} \cdot \epsilon \quad \text{and} \quad |\partial_d z_n|_1 \leq \frac{1}{2^n} \cdot \epsilon.$$

We now use UBC to construct a locally finite fundamental cycle of  $M$  out of these relative cycles: For  $n \in \mathbb{N}$ , the boundary parts  $C_{d-1}(i_n; \mathbb{R})(\partial_d z_n|_{B_{n,+}})$

and  $-C_{d-1}(i_{n+1}; \mathbb{R})(\partial_d z_{n+1}|_{B_{n+1,-}})$  are fundamental cycles of the sphere  $S^{d-1}$  (embedded via  $i_n \circ i_{n,+}$  and  $i_{n+1} \circ i_{n+1,-}$  into  $M$ , which implicitly uses the orientation-reversing homeomorphism  $f_n$ ). By UBC, there exists a chain  $b_n \in C_d(S^{d-1}; \mathbb{R})$  with

$$\begin{aligned} \partial_d C_d(i_n \circ i_{n,+}; \mathbb{R})(b_n) &= C_{d-1}(i_n; \mathbb{R})(\partial_d z_n|_{B_{n,+}}) \\ &\quad + C_{d-1}(i_{n+1}; \mathbb{R})(\partial_d z_{n+1}|_{B_{n+1,-}}) \end{aligned}$$

and

$$|b_n|_1 \leq K \cdot \left( \frac{1}{2^n} + \frac{1}{2^{n+1}} \right) \cdot \epsilon \leq K \cdot \frac{1}{2^{n-1}} \cdot \epsilon.$$

A straightforward computation shows that

$$c := \sum_{n=0}^{\infty} C_d(i_n; \mathbb{R})(z_n - C_d(i_{n,+}; \mathbb{R})(b_n))$$

is a locally finite  $d$ -cycle on  $M$ . Moreover, the local contribution on  $W_0$  shows that  $c$  is a locally finite fundamental cycle of  $M$ . By construction,

$$\begin{aligned} |c|_1 &\leq \sum_{n=0}^{\infty} (|z_n|_1 + |b_n|_1) \\ &\leq \sum_{n=0}^{\infty} \left( \alpha_n + \frac{1}{2^n} \cdot \epsilon + K \cdot \frac{1}{2^{n-1}} \cdot \epsilon \right) \leq \sum_{n=0}^{\infty} \alpha_n + (2 + 4 \cdot K) \cdot \epsilon \\ &= \alpha + (2 + 4 \cdot K) \cdot \epsilon. \end{aligned}$$

Thus, taking  $\epsilon \rightarrow 0$ , we obtain  $\|M\|^{\text{lf}} \leq \alpha$ .  $\square$

**Claim 2.2.** *We have  $\|M\|^{\text{lf}} \geq \alpha$ .*

*Proof.* Without loss of generality we may assume that  $\|M\|^{\text{lf}}$  is finite. Let  $c \in C_d^{\text{lf}}(M; \mathbb{R})$  be a locally finite fundamental cycle of  $M$  with  $|c|_1 < \infty$ . For  $n \in \mathbb{N}$ , we consider the subchain  $c_n := c|_{W_{(n)}}$  of  $c$ , consisting of all simplices whose images touch  $W_{(n)} := \bigcup_{k=0}^n i_k(W_k) \subset M$ . Because  $c$  is locally finite, each  $c_n$  is a finite singular chain and  $(|c_n|_1)_{n \in \mathbb{N}}$  is a monotonically increasing sequence with limit  $|c|_1$ .

Let  $\epsilon \in \mathbb{R}_{>0}$ . Then there is an  $n \in \mathbb{N}_{>0}$  that satisfies  $|c - c_n|_1 \leq \epsilon$  and  $\alpha - \sum_{k=0}^n \alpha_k \leq \epsilon$ . Let

$$p: M \rightarrow W_{(n)}/i_n(B_{n,+}) =: W$$

be the map that collapses everything beyond stage  $n+1$  to a single point  $x$ . Then  $z := C_d(p; \mathbb{R})(c_n) \in C_d(W, \{x\}; \mathbb{R})$  is a relative cycle and

$$|\partial_d z|_1 \leq |\partial_d c_n|_1 \leq |\partial_d(c - c_n)|_1 \leq (d+1) \cdot |c - c_n|_1 \leq (d+1) \cdot \epsilon.$$

Because  $d > 1$ , there exists a chain  $b \in C_d(\{x\}; \mathbb{R})$  with

$$\partial_d b = \partial_d z \quad \text{and} \quad |b|_1 \leq |\partial_d z|_1 \leq (d+1) \cdot \epsilon.$$

Then

$$\bar{z} := z - b \in C_d(W; \mathbb{R})$$

is a cycle on  $W$ ; because  $z$  and  $\bar{z}$  have the same local contribution on  $W_0$ , the cycle  $z$  is a fundamental cycle of the manifold

$$W \cong M_0 \# \cdots \# M_n.$$

As  $d > 2$ , the construction of our chains and additivity of simplicial volume under connected sums [Gro82, BBF<sup>+</sup>14] show that

$$\begin{aligned} |c|_1 &\geq |c_n|_1 \geq |z|_1 \geq |\bar{z}|_1 - |b|_1 \\ &\geq \|W\| - (d+1) \cdot \epsilon = \sum_{k=0}^n \|M_k\| - (d+1) \cdot \epsilon \\ &\geq \alpha - (d+2) \cdot \epsilon. \end{aligned}$$

Thus, taking  $\epsilon \rightarrow 0$ , we obtain  $|c|_1 \geq \alpha$ ; hence,  $\|M\|^{\text{lf}} \geq \alpha$ .  $\square$

This completes the proof of Theorem A.

*Remark 2.3* (adding geometric structures). In fact, this argument can also be performed smoothly: The constructions leading to Theorem 1.2 can be carried out in the smooth setting. Therefore, we can choose the  $(M_n)_{n \in \mathbb{N}}$  to be smooth and equip  $M$  with a corresponding smooth structure. Moreover, we can endow these smooth pieces with Riemannian metrics. Scaling these Riemannian metrics appropriately shows that we can turn  $M$  into a Riemannian manifold of finite volume.

### 3 Proof of Theorem B

In this section, we prove Theorem B, i.e., that the set of simplicial volumes of tame manifolds is countable.

**Definition 3.1.** A manifold  $M$  without boundary is *tame* if there exists a compact connected manifold  $W$  with boundary such that  $M$  is homeomorphic to  $W^\circ := W \setminus \partial W$ .

As in the closed case, our proof is based on a counting argument:

**Proposition 3.2.** *There are only countably many proper homotopy types of tame manifolds.*

As we could not find a proof of this statement in the literature, we will give a complete proof in Section 3.1 below. Theorem B is a direct consequence of Proposition 3.2:

*Proof of Theorem B.* The simplicial volume  $\|\cdot\|^{\text{lf}}$  is invariant under proper homotopy equivalence (this can be shown as in the compact case). Therefore, the countability of  $\text{SV}^{\text{lf}}(d)$  follows from the countability of the set of proper homotopy types of tame  $d$ -manifolds (Proposition 3.2).  $\square$

*Remark 3.3.* Let  $d \in \mathbb{N}_{\geq 3}$ . Then  $\infty \in \text{SV}_{\text{tame}}^{\text{lf}}(d)$ : Let  $N$  be an oriented closed connected hyperbolic  $(d-1)$ -manifold and let  $M := N \times \mathbb{R}$ . Then  $M$  is tame (as interior of  $N \times [0, 1]$ ) and  $\|N\| > 0$  [Gro82, Section 0.3][Thu97, Theorem 6.2]. Hence, by the finiteness criterion [Gro82, p. 17][Löh08, Theorem 6.4], we obtain that  $\|M\|^{\text{lf}} = \infty$ .

### 3.1 Counting tame manifolds

It remains to prove Proposition 3.2. We use the following observations:

**Definition 3.4** (models of tame manifolds).

- A *model* of a tame manifold  $M$  is a finite CW-pair  $(X, A)$  (i.e., a finite CW-complex  $X$  with a finite subcomplex  $A$ ) that is homotopy equivalent (as pairs of spaces) to  $(W, \partial W)$ , where  $W$  is a compact connected manifold with boundary whose interior is homeomorphic to  $M$ .
- Two models of tame manifolds are *equivalent* if they are homotopy equivalent as pairs of spaces.

**Lemma 3.5** (existence of models). *Let  $W$  be a compact connected manifold. Then there exists a finite CW-pair  $(X, A)$  such that  $(W, \partial W)$  and  $(X, A)$  are homotopy equivalent pairs of spaces.*

*In particular: Every tame manifold admits a model.*

*Proof.* It should be noted that we work with topological manifolds; hence, we cannot argue directly via triangulations. Of course, the main ingredient is the fact that every compact manifold is homotopy equivalent to a finite complex [Sie68, KS69].

Hence, there exist finite CW-complexes  $A$  and  $Y$  with homotopy equivalences  $f: A \rightarrow \partial W$  and  $g: Y \rightarrow W$ . Let  $j := \bar{g} \circ i \circ f$ , where  $i: \partial W \hookrightarrow W$  is the inclusion and  $\bar{g}$  is a homotopy inverse of  $g$ . By construction, the upper square in the diagram in Figure 2 is homotopy commutative.

As next step, we replace  $j: A \rightarrow Y$  by a homotopic map  $j_c: A \rightarrow Y$  that is cellular (second square in Figure 2).

The mapping cylinder  $Z$  of  $j_c$  has a finite CW-structure (as  $j_c$  is cellular) and the canonical map  $p: Z \rightarrow Y$  allows to factor  $j_c$  into an inclusion  $J$  of a subcomplex and the homotopy equivalence  $p$  (third square in Figure 2).

We thus obtain a homotopy commutative square

$$\begin{array}{ccc} \partial W & \xrightarrow{i} & W \\ f \uparrow & \text{\textcircled{+}} & \uparrow F := g \circ p \\ A & \xrightarrow{J} & Z \end{array}$$

where the vertical arrows are homotopy equivalences, the upper horizontal arrow is the inclusion, and the lower horizontal arrow is the inclusion of a subcomplex.

Using a homotopy between  $i \circ f$  and  $F \circ J$  and adding another cylinder to  $Z$ , we can replace  $Z$  by a finite CW-complex  $X$  (that still contains  $A$  as subcomplex) to obtain a *strictly* commutative diagram

$$\begin{array}{ccc} \partial W & \xrightarrow{i} & W \\ f \uparrow \simeq & & \simeq \uparrow \\ A & \longrightarrow & X \end{array}$$

whose vertical arrows are homotopy equivalences and whose horizontal arrows are inclusions.

$$\begin{array}{ccc}
\partial W & \xrightarrow{i} & W \\
f \uparrow & \text{\scriptsize } \circlearrowleft h \text{\scriptsize } & g \uparrow \downarrow \bar{g} \\
A & \xrightarrow{j} & Y \\
\parallel & \text{\scriptsize } \circlearrowleft h \text{\scriptsize } & \parallel \\
A & \xrightarrow{j_c} & Y \\
\parallel & & \parallel \\
A & \xrightarrow{J} & Z \\
& & \uparrow p
\end{array}$$

Figure 2: Finding a model

Because the inclusions  $\partial W \hookrightarrow W$  (as inclusion of the boundary of a compact topological manifold) and  $A \hookrightarrow X$  (as inclusion of a subcomplex) are cofibrations, this already implies that the vertical arrows form a homotopy equivalence  $(X, A) \rightarrow (W, \partial W)$  of pairs [May99, Chapter 6.5].  $\square$

**Lemma 3.6** (equivalence of models). *If  $M$  and  $N$  are tame manifolds with equivalent models, then  $M$  and  $N$  are properly homotopy equivalent.*

*Proof.* As  $M$  and  $N$  admit equivalent models, there exist compact connected manifolds  $W$  and  $V$  with boundary such that  $M \cong W^\circ$  and  $N \cong V^\circ$  and such that the pairs  $(W, \partial W)$  and  $(V, \partial V)$  are homotopy equivalent (by transitivity of homotopy equivalence of pairs of spaces). Let  $(f, f_\partial): (W, \partial W) \rightarrow (V, \partial V)$  and  $(g, g_\partial): (V, \partial V) \rightarrow (W, \partial W)$  be mutually homotopy inverse homotopy equivalences of pairs.

By the topological collar theorem [Bro62, Con71], we have homeomorphisms

$$\begin{aligned}
M &\cong W \cup_{\partial W} (\partial W \times [0, \infty)) \\
N &\cong V \cup_{\partial V} (\partial V \times [0, \infty)),
\end{aligned}$$

where the glueing occurs via the canonical inclusions  $\partial W \hookrightarrow \partial W \times [0, \infty)$  and  $\partial V \hookrightarrow \partial V \times [0, \infty)$  at parameter 0.

Then the maps  $f$  and  $f_\partial \times \text{id}_{[0, \infty)}$  glue to a well-defined proper continuous map  $F: M \rightarrow N$  and the maps  $g$  and  $g_\partial \times \text{id}_{[0, \infty)}$  glue to a well-defined proper continuous map  $G: N \rightarrow M$ .

Moreover, the homotopy of pairs between  $(f \circ g, f_\partial \circ g_\partial)$  and  $(\text{id}_V, \text{id}_{\partial V})$  glues into a proper homotopy between  $F \circ G$  and  $\text{id}_M$ . In the same way, there is a proper homotopy between  $G \circ F$  and  $\text{id}_N$ . Hence, the spaces  $M$  and  $N$  are properly homotopy equivalent.  $\square$

**Lemma 3.7** (countability of models). *There exist only countably many equivalence classes of models.*

*Proof.* There are only countably many homotopy types of finite CW-complexes (because every finite CW-complex is homotopy equivalent to a finite simplicial

complex). Moreover, every finite CW-complex has only finitely many subcomplexes. Therefore, there are only countably many homotopy types (of pairs of spaces) of finite CW-pairs.  $\square$

*Proof of Proposition 3.2.* We only need to combine Lemma 3.5, Lemma 3.6, and Lemma 3.7.  $\square$

## 4 Low dimensions

### 4.1 Dimension 2

We now compute the set of simplicial volumes of surfaces. We first consider the tame case:

**Example 4.1** (tame surfaces). Let  $W$  be an oriented compact connected surface with  $g \in \mathbb{N}$  handles and  $b \in \mathbb{N}$  boundary components. Then the proportionality principle for simplicial volume of hyperbolic manifolds [Gro82, p. 11] (a thorough exposition is given, for instance, by Fujiwara and Manning [FM11, Appendix A]) gives

$$\|W^\circ\|^{\text{lf}} = \begin{cases} 4 \cdot (g - 1) + 2 \cdot b & \text{if } g > 0 \\ 2 \cdot b - 4 & \text{if } g = 0 \text{ and } b > 1 \\ 0 & \text{if } g = 0 \text{ and } b \in \{0, 1\}. \end{cases}$$

**Proposition 4.2.** *We have  $\text{SV}^{\text{lf}}(2) = 2 \cdot \mathbb{N} \cup \{\infty\}$  and  $\text{SV}_{\text{tame}}^{\text{lf}}(2) = 2 \cdot \mathbb{N}$ .*

*Proof.* We first prove  $2 \cdot \mathbb{N} \subset \text{SV}_{\text{tame}}^{\text{lf}}(2) \subset \text{SV}^{\text{lf}}(2)$  and  $\infty \in \text{SV}^{\text{lf}}(2)$ , i.e., that all the given values may be realised: In view of Example 4.1, all even numbers occur as simplicial volume of some (possibly open) tame surface.

Let

$$M := T^2 \# T^2 \# T^2 \# \dots$$

be an infinite “linear” connected sum of tori  $T^2$ . Collapsing  $M$  to the first  $g \in \mathbb{N}$  summands and an argument as in the proof of Claim 2.2 shows that

$$\|M\|^{\text{lf}} \geq \|\Sigma_g\| = 4 \cdot g - 4$$

for all  $g \in \mathbb{N}_{\geq 1}$ . Hence,  $\|M\|^{\text{lf}} = \infty$ .

It remains to show that  $\text{SV}^{\text{lf}}(2) \subset 2 \cdot \mathbb{N} \cup \{\infty\}$ : Let  $M$  be an oriented connected (topological, separable, Hausdorff) 2-manifold without boundary. Then  $M$  admits a smooth structure [Moi77] and whence a proper smooth map  $p: M \rightarrow \mathbb{R}$ . Using suitable regular values of  $p$ , we can thus write  $M$  as an ascending union

$$M = \bigcup_{n \in \mathbb{N}} M_n$$

of oriented connected compact submanifolds (possibly with boundary)  $M_n$  that are nested via  $M_0 \subset M_1 \subset \dots$ . Then one of the following cases occurs:

1. There exists an  $N \in \mathbb{N}$  such that for all  $n \in \mathbb{N}_{\geq N}$  the inclusion  $M_n \hookrightarrow M_{n+1}$  is a homotopy equivalence.
2. For each  $N \in \mathbb{N}$  there exists an  $n \in \mathbb{N}_{\geq N}$  such that the inclusion  $M_n \hookrightarrow M_{n+1}$  is *not* a homotopy equivalence.

In the first case, the classification of compact surfaces with boundary shows that  $M$  is tame. Hence  $\|M\|^{\text{lf}} \in 2 \cdot \mathbb{N}$  (Example 4.1).

In the second case, the manifold  $M$  is *not* tame (which can, e.g., be derived from the classification of compact surfaces with boundary). We show that  $\|M\|^{\text{lf}} = \infty$ . To this end, we distinguish two cases:

- a. The sequence  $(h(M_n))_{n \in \mathbb{N}}$  is unbounded, where  $h(\cdot)$  denotes the number of handles of the surface.
- b. The sequence  $(h(M_n))_{n \in \mathbb{N}}$  is bounded.

In the unbounded case, a collapsing argument (similar to the argument for  $T^2 \# T^2 \# \dots$  and Claim 2.2) shows that  $\|M\|^{\text{lf}} = \infty$ .

We claim that also in the bounded case we have  $\|M\|^{\text{lf}} = \infty$ : Shifting the sequence in such a way that all handles are collected in  $M_0$ , we may assume without loss of generality that the sequence  $(h(M_n))_{n \in \mathbb{N}}$  is constant. Thus, for each  $n \in \mathbb{N}$ , the surface  $M_{n+1}$  is obtained from  $M_n$  by adding a finite disjoint union of disks and of spheres with finitely many (at least two) disks removed; we can reorganise this sequence in such a way that no disks are added. Hence, we may assume that  $M_n$  is a retract of  $M_{n+1}$  for each  $n \in \mathbb{N}$ . Furthermore, because we are in case 2, the classification of compact surfaces shows (with the help of Example 4.1) that

$$\lim_{n \rightarrow \infty} \|M_n\| = \infty.$$

Let  $c \in C_2^{\text{lf}}(M; \mathbb{R})$  be a locally finite fundamental cycle of  $M$  and let  $n \in \mathbb{N}$ . Because  $c$  is locally finite, there is a  $k \in \mathbb{N}$  such that  $c|_{M_n}$  is supported on  $M_{n+k}$ ; the restriction  $c|_{M_n}$  consists of all summands of  $c$  whose supports intersect with  $M_n$ . Because  $M_n$  is a retract of  $M_{n+k}$ , we obtain from  $c|_{M_n}$  a relative fundamental cycle  $c_n$  of  $M_n$  by pushing the chain  $c|_{M_n}$  to  $M_n$  via a retraction  $M_{n+k} \rightarrow M_n$ . Therefore,

$$|c|_1 \geq |c|_{M_n}|_1 \geq |c_n|_1 \geq \|M_n\|.$$

Taking  $n \rightarrow \infty$  shows that  $|c|_1 = \infty$ . Taking the infimum over all locally finite fundamental cycles  $c$  of  $M$  proves that  $\|M\|^{\text{lf}} = \infty$ .

Moreover, Example 4.1 shows that  $\infty \notin \text{SV}_{\text{tame}}^{\text{lf}}(2)$ .  $\square$

## 4.2 Dimension 3

The general case of non-compact 3-manifolds seems to be rather involved (as the structure of non-compact 3-manifolds can get fairly complicated). We can at least deal with the tame case:

**Proposition 4.3.** *We have  $\text{SV}_{\text{tame}}^{\text{lf}}(3) = \text{SV}(3) \cup \{\infty\}$ .*

*Proof.* Clearly,  $\text{SV}(3) \subset \text{SV}_{\text{tame}}^{\text{lf}}(3)$  and  $\infty \in \text{SV}_{\text{tame}}^{\text{lf}}(3)$  (Remark 3.3).

Conversely, let  $W$  be an oriented compact connected 3-manifold and let  $M := W^\circ$ . We distinguish the following cases:

- If at least one of the boundary components of  $W$  has genus at least 2, then the finiteness criterion [Gro82, p. 17][Löh08, Theorem 6.4] shows that  $\|M\|^{\text{lf}} = \infty$ .

- If the boundary of  $W$  consists only of spheres and tori, then we proceed as follows: In a first step, we fill in all spherical boundary components of  $W$  by 3-balls and thus obtain an oriented compact connected 3-manifold  $V$  all of whose boundary components are tori. In view of considerations on tame manifolds with amenable boundary [KK15] and glueing results for bounded cohomology [Gro82][BBF<sup>+</sup>14], we obtain that

$$\|M\|^{\text{lf}} = \|W\| = \|V\|.$$

By Kneser's prime decomposition theorem [AFW15, Theorem 1.2.1] and the additivity of (relative) simplicial volume with respect to connected sums [Gro82][BBF<sup>+</sup>14] in dimension 3, we may assume that  $V$  is prime (i.e., admits no non-trivial decomposition as a connected sum). Moreover, because  $\|S^1 \times S^2\| = 0$ , we may even assume that  $V$  is irreducible [AFW15, p. 3].

By geometrisation [AFW15, Theorem 1.7.6], then  $V$  admits a decomposition along finitely many incompressible tori into Seifert fibred manifolds (which have trivial simplicial volume [Thu97, Corollary 6.5.3]) and hyperbolic pieces  $V_1, \dots, V_k$ . As the tori are incompressible, we can now again apply additivity [Gro82][BBF<sup>+</sup>14] to conclude that

$$\|V\| = \sum_{j=1}^k \|V_j\|.$$

Let  $j \in \{1, \dots, k\}$ . Then the boundary components of  $V_j$  are  $\pi_1$ -injective tori (as the interior of  $V_j$  admits a complete hyperbolic metric of finite volume) [BP92, Proposition D.3.18]. Let  $S$  be a Seifert 3-manifold whose boundary is a  $\pi_1$ -injective torus (e.g., the knot complement of a non-trivial torus knot [Mos71, Theorem 2][Lüc02, Lemma 4.4]). Filling each boundary component of  $V_j$  with a copy of  $S$  results in an oriented closed connected 3-manifold  $N_j$ , which satisfies (again, by additivity)

$$\|N_j\| = \|V_j\| + 0 = \|V_j\|.$$

Therefore, the oriented closed connected 3-manifold  $N := N_1 \# \dots \# N_k$  satisfies

$$\|N\| = \sum_{j=1}^k \|N_j\| = \sum_{j=1}^k \|V_j\| = \|V\|.$$

In particular,  $\|M\|^{\text{lf}} = \|V\| = \|N\| \in \text{SV}(3)$ . □

## References

- [AFW15] Matthias Aschenbrenner, Stefan Friedl, and Henry Wilton. *3-manifold groups*. EMS Series of Lectures in Mathematics. European Mathematical Society (EMS), Zürich, 2015. Cited on page: 10
- [BBF<sup>+</sup>14] M. Bucher, M. Burger, R. Frigerio, A. Iozzi, C. Pagliantini, and M. B. Pozzetti. Isometric embeddings in bounded cohomology. *J. Topol. Anal.*, 6(1):1–25, 2014. Cited on page: 3, 5, 10

- [BKK14] Michelle Bucher, In Kang Kim, and Sungwoon Kim. Proportionality principle for the simplicial volume of families of  $\mathbb{Q}$ -rank 1 locally symmetric spaces. *Math. Z.*, 276(1-2):153–172, 2014. Cited on page: 1
- [BP92] Riccardo Benedetti and Carlo Petronio. *Lectures on Hyperbolic Geometry*. Universitext. Springer-Verlag, Berlin, 1992. Cited on page: 10
- [Bro62] Morton Brown. Locally flat imbeddings of topological manifolds. *Ann. of Math. (2)*, 75:331–341, 1962. Cited on page: 7
- [Con71] Robert Connelly. A new proof of Brown’s collaring theorem. *Proc. Amer. Math. Soc.*, 27:180–182, 1971. Cited on page: 7
- [FM11] Koji Fujiwara and Jason Fox Manning. Simplicial volume and fillings of hyperbolic manifolds. *Algebr. Geom. Topol.*, 11(4):2237–2264, 2011. Cited on page: 8
- [FM18] Roberto Frigerio and Marco Moraschini. Gromov’s theory of multicomplexes with applications to bounded cohomology and simplicial volume. *Mem. Amer. Math. Soc.*, to appear, 2018. arXiv:1808.07307 [math.GT]. Cited on page: 1
- [Gro82] Michael Gromov. Volume and bounded cohomology. *Inst. Hautes Études Sci. Publ. Math.*, (56):5–99 (1983), 1982. Cited on page: 1, 3, 5, 8, 9, 10
- [HL19] Nicolaus Heuer and Clara Löh. Transcendental simplicial volumes. 2019. preprint. Cited on page: 2
- [HL20] Nicolaus Heuer and Clara Löh. The spectrum of simplicial volume. *Invent. math.*, 2020. DOI 10.1007/s00222-020-00989-0. Cited on page: 1
- [KK15] Sungwoon Kim and Thilo Kuessner. Simplicial volume of compact manifolds with amenable boundary. *J. Topol. Anal.*, 7(1):23–46, 2015. Cited on page: 1, 10
- [KS69] R. C. Kirby and L. C. Siebenmann. On the triangulation of manifolds and the Hauptvermutung. *Bull. Amer. Math. Soc.*, 75:742–749, 1969. Cited on page: 6
- [Löh08] Clara Löh. Isomorphisms in  $l^1$ -homology. *Münster J. Math.*, 1:237–265, 2008. Cited on page: 1, 5, 9
- [LS09a] Clara Löh and Roman Sauer. Degree theorems and Lipschitz simplicial volume for nonpositively curved manifolds of finite volume. *J. Topol.*, 2(1):193–225, 2009. Cited on page: 1
- [LS09b] Clara Löh and Roman Sauer. Simplicial volume of Hilbert modular varieties. *Comment. Math. Helv.*, 84(3):457–470, 2009. Cited on page: 1
- [Lüc02] Wolfgang Lück.  $L^2$ -Invariants: Theory and Applications to Geometry and K-Theory, volume 44 of *Ergebnisse der Mathematik und ihrer Grenzgebiete. 3. Folge. A Series of Modern Surveys in Mathematics [Results in Mathematics and Related Areas. 3rd Series. A Series of Modern Surveys in Mathematics]*. Springer-Verlag, Berlin, 2002. Cited on page: 10
- [May99] J. Peter May. *A concise course in algebraic topology*. Chicago Lectures in Mathematics. University of Chicago Press, Chicago, IL, 1999. Cited on page: 7
- [MM85] Shigenori Matsumoto and Shigeyuki Morita. Bounded cohomology of certain groups of homeomorphisms. *Proc. Amer. Math. Soc.*, 94(3):539–544, 1985. Cited on page: 3
- [Moi77] Edwin E. Moise. *Geometric topology in dimensions 2 and 3*, volume 47 of *Graduate Texts in Mathematics*. Springer-Verlag, New York-Heidelberg, 1977. Cited on page: 8

- [Mos71] Louise Moser. Elementary surgery along a torus knot. *Pacific J. Math.*, 38:737–745, 1971. Cited on page: 10
- [Sie68] L. C. Siebenmann. On the homotopy type of compact topological manifolds. *Bull. Amer. Math. Soc.*, 74:738–742, 1968. Cited on page: 6
- [Thu97] William P. Thurston. *Three-dimensional geometry and topology. Vol. 1*, volume 35 of *Princeton Mathematical Series*. Princeton University Press, Princeton, NJ, 1997. Edited by Silvio Levy. Cited on page: 5, 10

*Nicolaus Heuer*

DPMMS, University of Cambridge  
nh441@cam.ac.uk, <https://www.dpmms.cam.ac.uk/~nh441>

*Clara Löh*

Fakultät für Mathematik, Universität Regensburg, 93040 Regensburg  
clara.loeh@mathematik.uni-r.de, <http://www.mathematik.uni-r.de/loeh>